Assessment Overview

Assessment Area

ICF Domain: Body Function Subcategory: Mental Functions

You Will Need

Length: 5 minutes, 28 items, selfadministered questionnaire. Scoring: Respondents rate each question using options provided: 1) "Better than usual" 2) "Same as usual" 3) "Worse than usual" 4) "Much worse than usual" Different methods of scoring are possible, which will affect the total score. Traditional scoring

method assigns score of 0 for responses 1 and 2 and a score of 1 for responses 3 and 4.

Summary

The General Health Questionnaire-28 (GHQ-28) is a self-report questionnaire used to indicate psychological well-being and detect possible cases of psychiatric distress related to general medical illness.

The GHQ-28 is designed to assess 4 aspects of distress: 1) depression, 2) anxiety, 3) social impairment, 4) hypochondriasis

Availability

Worksheet: Can be purchased here.

Languages: Available in 38 languages

Assessment Interpretability

Minimal Clinically Important Difference	Statistical Error	Typical Values
Not established in SCI	Not established in SCI	Not established in SCI

Measurement Properti	es	
Validity – High		Reliability – High
High correlation with Clinica	l Interview Scale:	High Internal Consistency:
r=0.83		Cronbach's α = 0.82
(Griffiths et al. 1993; n=60, 77% males; duration of paralysis: 7 years)	median age: 33.8 years; median	(Griffiths et al. 1993; n=60, 77% males; median age: 33.8 years; median duration of paralysis: 7 years)
Number of studies reporting	s validity data: 1	Number of studies reporting reliability data: $f 1$
	Respo	nsiveness
Floor/Ceiling Effect:	Effect Size:	Number of studies reporting

Floor/Ceiling Effect: Not established in SCI Effect Size: Not established in SCI Number of studies reporting responsiveness data: 0