

Braden Scale

Assessment Overview

Assessment Area

ICF Domain:

Body Function

Subcategory:

Functions of the Skin

Subscales (domains):

- 1) Sensory Perception, 2) Moisture, 3) Activity, 4) Mobility, 5) Nutrition, 6) Friction and Shear

You Will Need

Length:

5-10 minutes, 6 items

Scoring:

Each domain scored 1-4 (except for Friction and Shear, which is scored 1-3), total score (6-23) as sum of domains.

Higher scores reflect better prognosis.

Summary

The Braden Scale is a clinician-administered assessment tool for determining a patient's risk level for incurring skin breakdown and is useful for detecting pressure ulcer risk in people with SCI (though it includes two factors less related to risk for people with SCI - sensory perception and mobility). Moisture was found to be the most predictive variable for people with SCI. It has been tested in both acute care and long-term-care settings.

The scale items were developed based on expert consensus, and includes three factors (sensory perception, mobility and nutritional variables) that were not significantly related to pressure ulcer development for individuals with SCI.

Availability

Available for free here: www.bradenscale.com

Assessment form available here:

http://www.in.gov/isdh/files/Braden_Scale.pdf

Languages: English, French, Portuguese / Brazilian Portuguese, Spanish. Also available in other languages, but are not formally validated.

Video: <https://www.youtube.com/watch?v=zP6p0M5zrLk&t=10s>

Assessment Interpretability

Minimal Clinically Important Difference

Not established in SCI

Statistical Error

Not established in SCI

Typical Values

Mean (95%CI) Scores:

All patients: 11.1 (10.7-11.5)

Patients with ulcers (n=80): 9.9 (9.6-10.3)

Patients without ulcers (n=64): 12.6 (12-13.2)

(Ash 2002; n=144; mixed injury types; mean time since injury = 14 days)

Mean (SD) Scores:

13.8 (1.75) (range 10-18)

(Wellard, 2002; N=60; mixed injury types; non-acute SCI patients)

Threshold Values:

16 or less indicates risk of pressure ulcer

(Flett et al. 2019; n=754 (510 males); mean age (SD): 53.9 (18.5); Tetraplegic 43%, Paraplegic 7%; Complete injury 15%, Incomplete 77%; injury duration (SD): 84.6 (378.4) days)

Measurement Properties

Validity – **Moderate**

Moderate correlation with the stage of the first pressure ulcer:

r = -0.353

Moderate correlation with the number of ulcers developed:

r = -0.431

(Salzberg et al. 1999; n=226, 188 males; mixed injury types; acute, traumatic SCI)

Moderate predictive validity for pressure ulcer development:

Area Under Curve (AUC) = 0.73-0.81

CI (95%) = 0.74-0.88

(Ash 2002; n=144; mixed injury types; mean time since injury = 14 days)

Study findings suggest that a simple measure of mobility, admission FIM bed/chair transfer score of 1 (total assist), can identify at-risk individuals with greater accuracy than both an SCI specific instrument (SCIPUS) and a PI specific instrument (Braden)

(Flett et al. 2019; n=754 (510 males); mean age (SD): 53.9 (18.5); Tetraplegic 43%, Paraplegic 7%; Complete injury 15%, Incomplete 77%; injury duration (SD): 84.6 (378.4) days)

It was found that sensory perception, mobility and nutritional variables were not significantly related to pressure ulcer development. Moisture was the most important predictive variable

(Salzberg et al., 1999; N=226, 188 male; acute traumatic SCI, mixed injury types)

Number of studies reporting validity data: 4

Reliability

Not established in SCI

Responsiveness

Floor/Ceiling Effect:

A ceiling effect was reported in mixed populations (21% of patients attained a 'high risk' score)

(Wellard, 2002; N=60; mixed injury types; non-acute SCI patients)

Effect Size:

Not established in SCI

Number of studies reporting responsiveness data: 2