# **Assessment Overview**

# Assessment Area

**ICF Domain:** 

Activity

Subcategory:

Self-Care

# You Will Need

### Length:

13 items – less than 15 minutes

## Scoring:

4 items (washing up, washing clothes, driving a car/bus travel, and gainful work) are scored on a 2-point scale; the remaining 9 items are scored on a 3-point scale.

Response categories are 0 = never, 1 = occasionally or more, and 2 = most days.

# Summary

The Frenchay Activities Index assesses frequency of performing Instrumental Activities of Daily Living, including items that reflect the patient's behavior in areas of domestic chores, leisure/work and outdoor activities.

The raw scores of the 13-item FAI can be transformed to interval scores, thereby enabling clinicians and researchers to quantify and monitor IADL function in SCI patients.

# Availability

Available for free here:

http://scire-nexcess.developmentwebsite.ca/wp-

# **Assessment Interpretability**

# Minimal Clinically Important Difference

Not established in SCI

# Statistical Error

#### Standard Error (SE) of the items:

1. Washing up	0.17
2. Preparing main meals	0.13
3. Washing clothes	0.17
<ol><li>Driving car/bus travel</li></ol>	0.17
5. Light housework	0.11
<ol><li>Heavy housework</li></ol>	0.12
<ol><li>Local shopping</li></ol>	0.11
8. Social occasions	0.13
<ol><li>Actively pursuing hobby</li></ol>	0.12
10. Gainful work	0.16
11. Travel outings/car rides	0.15
12. Gardening	0.15
13. Household/car maintenance	0.15

(Hsieh et al. 2007)

# Typical Values

Not established in SCI

# Validity – Low to Moderate

<u>Low</u> to <u>Moderate</u> Correlation between R-FAI administered at 3, 6, and 12 months after injury and the 4 domains of the WHOQOL- BREF administered at 12 months after injury:

WHOQOL-			
BREF 12			
months post	FAI 3 months	FAI 6 months	FAI 12 months
injury	post injury	post injury	post injury
Physical	r = 0.39	r = 0.41	r = 0.50
Psychology	r = 0.38	r = 0.28	r = 0.37
Social relations	r = 0.20	r = 0.28	r = 0.35
Environment	r = 0.39	r = 0.31	r = 0.37

(Chern et al. 2013, N=2339 (1454 male), mean age: 45 (SD 18.5), traumatic limb injuries)

<u>Not Ranked</u> The Frenchay Activities Index was validated as a unidimensional construct through revision of the scale after Rasch analysis.

(Hsieh et al. 2007, N=233 (193 male), mean age: 41.1 (SD 12.6), 33 Complete, 57 Incomplete tetraplegia; 151 Complete, 48 Incomplete paraplegia)

Number of studies reporting validity data: 2

# Reliability – High

# High Internal Consistency:

 $\alpha = >0.90$ 

(Chern et al. 2013, N=2339 (1454 male), mean age: 45 (SD 18.5), traumatic limb injuries)

## Not Ranked Rasch analysis reliability coefficient:

Rasch coefficient = 0.78

(Hsieh et al. 2007, N=233 (193 male), mean age: 41.1 (SD 12.6), 33 Complete, 57 Incomplete tetraplegia; 151 Complete, 48 Incomplete paraplegia)

Number of studies reporting reliability data: 2

# Responsiveness

## Floor/Ceiling Effect:

# months post injury	% Ceiling	% Floor
3	0.3	7.3
6	3.5	4.3
12	2.5	2.4

(Chern et al. 2013, N=2339 (1454 male), mean age: 45 (SD 18.5), traumatic limb injuries)

## **Effect Size:**

# months post injury	Effect size
3	0.10
6	0.35
12	0.15

(Chern et al. 2013, N=2339 (1454 male), mean age: 45 (SD 18.5), traumatic limb injuries)

# Number of studies reporting responsiveness data: (1)

(Chern et al. 2013, N=2339 (1454 male), mean age: 45 (SD 18.5), traumatic limb injuries)

# months post injury	Standardized Response Mean
3	0.20
6	0.52
12	0.23