

Spinal Cord Injury Pressure Ulcer Scale (SCIPUS)

Assessment Overview

Assessment Area

ICF Domain:

Body Function

Subcategory:

Functions of the Skin

You Will Need

Length:

15 items

Scoring:

By adding domain scores together a summary score (0-25) is calculated. Lower scores indicate better prognosis.

Summary

The Spinal Cord Injury Pressure Ulcer Scale (SCIPUS) is a clinician-administered measure developed to assess the risk for pressure ulcer development for individuals with SCI who are in a rehabilitation centre. Rating is based on personal knowledge of the client or chart review. Blood tests are required if tests for diabetes, albumin and hemocrit are not already part of the patient's medical record.

Availability

Available for free here:

<https://www.sralab.org/sites/default/files/2017-06/SCIPUS.pdf>

Languages: English

Assessment Interpretability

Minimal Clinically Important Difference

Not established in SCI

Statistical Error

Not established in SCI

Typical Values

Threshold Values:

Score of ≥ 6 has been suggested to indicate risk for pressure ulcer development.

(Salzberg et al. 1996; n=219, 217 male, traumatic SCI, mean (SD) time from injury to last follow up = 17.2 (12.1) years)

Measurement Properties

Validity – **Low** to **Moderate**

Moderate correlation with stage of first pressure ulcer:

$\rho = 0.34$

Moderate correlation with number of pressure ulcers:

$\rho = 0.34$

(Salzberg et al., 1999; N=226, 188 male, acute traumatic SCI, within 30 days of admission)

Low accuracy of “high risk” categorization (cut-off scores of ≥ 6 , ≥ 7 , ≥ 8):

AUC < 0.70

(Delparte et al., 2015; N=759, 509 male, mixed injury types, mean (SD) time since injury: 84.9 (379.7) days)

Number of studies reporting validity data: 3

Reliability – **Low** to **High**

High Inter-rater Reliability:

ICC = 0.91

(Delparte et al., 2015; N=759, 509 male, mixed injury types, mean (SD) time since injury: 84.9 (379.7) days)

Low Internal Consistency:

Person Separation Index (PSI)=0.44

(Higgins et al. 2019; N=886, 59% male; median age [interquartile range] 56 [28]; etiology: 56% non-traumatic, 44% traumatic)

Number of studies reporting reliability data: 2

Responsiveness

Floor/Ceiling Effect:

Not established in SCI

Effect Size:

Not established in SCI

Number of studies reporting

responsiveness data: 0