

<b>Reviewer ID:</b> Nicole Elfring, Jeremy Mak, Joanne Chi			
<b>Type of Outcome Measure:</b> Scaled-General Health Questionnaire (GHQ-28)			<b>Total articles:</b> 1
<b>Author ID Year</b>	<b>Study Design</b>	<b>Setting</b>	<b>Population (sample size, age) and Group</b>
Griffiths et al. 1993	Not specified	West Midlands Spinal Unit	N=60 (77% males) Median age = 33.8 (interquartile range (IQR) 27-47) years  Median duration of paralysis = 7 (IQR: 4-15) years Median age at paralysis = 23 (IQR: 19-35) years
<b>1. RELIABILITY</b> – no data available			
<b>2. VALIDITY</b>			
<b>Author ID</b>	<b>Validity</b>		
Griffiths 1993	GHQ-28 & Clinical Interview Scale (CIS) $r = 0.83$ (95 CI: 0.70-0.93)  Optimum discrimination occurred near GHQ 3/4 (0011 scoring scheme) giving a specificity of 0.82, sensitivity of 0.81 and a misclassification rate of 18%.		
<b>3. RESPONSIVENESS</b> – no data available			
<b>4. FLOOR/CEILING EFFECT</b> – no data available			
<b>5. INTERPRETABILITY</b> – no data available			