## Six-Minute Arm Test (6-MAT)

### **Assessment Overview**

#### Assessment Area

#### **ICF Domain:**

# Body Function Subcategory:

Functions & Structures of the Cardiovascular, Haematological, Immunological and Respiratory Systems

#### You Will Need

#### Length: 6 minutes

## Equipment:

- arm ergometer
- heart rate monitor
- RPE scale

#### Scoring:

The power output (PO) is selected for each individual based on their manual muscle strength, ASIA motor score and physical activity level (see table below). The aim is to attain a steady heart rate of 60%-70% of age-predicted maximum heart rate or a rating of 11-15 on the Borg RPE scale. Clinicians should record heart rate during the final 30 seconds of the test and the RPE at the end.

#### Summary

The Six-Minute Arm Test (6-MAT) assesses cardiovascular fitness in people with SCI. It involves 6 minutes of sub-maximal arm ergometry at a constant power output. This single stage test is simple and quick to administer clinically.

#### Availability

https://scireproject.com/wp-content/uploads/2022/05/worksheet\_6-mat.pdf

## **Assessment Interpretability**

Minimal Clinically Important Difference	Statistical Error	Typical Values
Not established in SCI	Standard Error of Measurement: Heart rate = 7.12 beats/min (95% Cl, 0.75- 0.96) VO2 = 1.62 mL/kg/min (95% Cl, 0.58-0.92) Minimal Detectable Change: Heart rate = 19.74 beats/min VO2 = 4.49 mL/kg/min (Hol et al. 2007; n=30, 25 males, mixed injury types, chronic SCl, mean time since injury (SD)= 12.0 (9.8) years)	Mean (SD) Scores: Heart rate (beats/min) = 129 (29), range = 75-183 VO2 peak (mL/kg/min) = 18.6 (8.4), range = 6.5-38.1 (Hol et al. 2007; n=30, 25 males, mixed injury types, chronic SCI, mean time since injury (SD)= 12.0 (9.8) years)

## **Measurement Properties**

Validity – <b>High</b>	Reliability – <b>High</b>	
Highcorrelation between 6-MAT VO2 and VO2 Peak:r = 0.91-0.92(Totosy de Zepetnek et al. 2016; n=52, mixed injury types, traumatic SCI, mean time since injury (SD)= 13.0 (10) years)Highcorrelation between 6-MAT Power Output and VO2 Peak: r = 0.73	<ul> <li><u>High</u> Test-retest Reliability (1 week interval): Heart Rate: ICC = 0.90</li> <li>VO<sub>2</sub>: ICC = 0.81</li> <li>(Hol et al. 2007; n=30, 25 males, mixed injury types, chronic SCI, mean time since injury (SD)= 12.0 (9.8) years)</li> <li>Number of studies reporting reliability data: 1</li> </ul>	
High       correlation between 6-MAT Heart Rate and VO2         Peak:       r = 0.63         (Hol et al. 2007; n=30, 25 males, mixed injury types, chronic SCI, mean time         circle injury (SD)= 12.0 (0.8) upper)		
Number of studies reporting validity data: 2		

## Responsiveness

Floor/Ceiling Effect: Not established in SCI Effect Size: Not established in SCI Number of studies reporting responsiveness data: 0