

Research Summary - Walking Index for Spinal Cord Injury (WISCI) - Lower Limb and Walking

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>Sato et al. 2023</p> <p>Japan</p> <p>Validity Study</p> <p>Rehabilitation hospital</p>	<p>N = 30 (5F)</p> <p>Mean age: 63.8 ±10.7 years, Tetraplegia = 17</p> <p>6 AIS A, 0 AIS B, 8 AIS C, 16 AIS D.</p> <p>Time since injury 1142 ±1720.7 days</p>	<p>Moderate to high correlation coefficient between the trunk assessment scale for spinal cord injury (TASS) and the WISCI II (r=0.67 (0.41-0.83))</p> <p>Construct validity for WISCI II with trunk control test (TCT-SCI) was r= 0.42 (0.14-0.71)</p>		
<p>Sinovas-Alonso et al. 2023</p> <p>Spain</p> <p>Observational, cross-sectional</p> <p>Biomechanics and Technical Aids Unit of the National Hospital for</p>	<p>iSCI</p> <p>N= 35 (24M) Mean age: 35.5(17.2)</p> <p>Non-SCI</p> <p>N = 50 (19M) Mean age: 34.6 (15.2)</p>	<p>Self-selected WISCI II levels showed good correlation with the spinal cord injury gait deviation index (SCI-GDI) (r=0.521)</p> <p>Maximum WISCI II levels had no significant correlations with the SCI-GDI (p=0.013)</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
Paraplegics of Toledo, Spain				
Willi et al. 2023 Switzerland Multicenter-observational study	N=50 Age range: 18-79 (52.6 ±16.2 years) Tetraplegic = 24 Paraplegic = 26; 2 AIS A, 0 AIS B, 7 AIS C, 41 AIS D Years since injury = 6.11 ± 9.8	Construct validity: Moderate relationship with the 2MWT, r=0.571 (0.356-0.784)		
Kahn et al. 2020 USA	N= 12 (11M, 1F) Mean age: 55.41± 11.65 years (32-73) Chronic motor SCI 2 AIS C, 10 AIS D Level of injury: 7 cervical, 5 thoracic	Convergent validity: For the WISCI II with the functional gait assessment (FGA) was high (spearman's rho= 0.74, p=0.006)		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
	Years since injury = 1.7 to 29.7 (7.8 ± 7.8)			
Calhoun et al. 2017 USA Mixed methods	N=52 (22M, 30F) Age range: (2-17) Tetraplegic=14 Paraplegic=38 AIS: 3A, 3B, 9C, 16D, 21Unknown Neurological level: 5 C1-C4, 2 C5-C8, 24 T1-S5, 21 Unknown		Intra-rater reliability ICC=0.997, CI=0.995-0.998 Inter-rater reliability ICC=0.97, CI=0.95-0.98	
Scivoletto et al. 2014 Test-Retest analysis, calculation of reliability and smallest real difference (SRD)	N=33 (28M, 5F) Mean age: 44 years AIS: 33D 32 AIS-D, 1 AIS-C Injury level: 20 cervical, 8 thoracic, 5 lumbar		Intra-rater reliability =0.975-0.999 Maximum WISCI II entire group: ICC=0.996 Maximum WISCI II Tetraplegics (n=20): ICC=0.994	Responsiveness: No data available Floor/Ceiling Effect: No data available Interpretability SEM (WISCI II) for tetraplegics = 0.401 (N=20); for paraplegics

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
SCI unit of a rehabilitation hospital	Median time since SCI onset = 40 days Incomplete SCI, subacute and chronic		Maximum WISCI II Paraplegics (n=13): ICC=0.992	= 0.437 (N=13); for both groups = 0.318. MDC for tetraplegics = 1.147 (N=20); for paraplegics = 1.682 (N=13); for both groups = 0.883
Tamburella et al. 2014 Serial cross-sectional study	N=23 (14M) Mean age 48.27 SD = 15.94 Mean time since injury = 16.43 months, SD = 19.03		Intra-rater ICC = 0.95, p<0.005	Responsiveness: ES = 0.07 Floor/Ceiling Effect: No data available Interpretability: SEM = 0.73, MDC95 = 0.02, %MDC = 13.0
Ovechkin et al. 2013 USA Prospective cohort study	N = 11 (3F, 8M) Age: 48 ± 19 AIS A: 4 AIS C: 1 AIS D: 6	AIS: Spearman rho = 0.71 (p< 0.05) FIM motor score: Spearman rho = 0.69 (p< 0.01) SCIM total score: Spearman rho = 0.74 (p<0.01)		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
University of Louisville		SCIM mobility score: Spearman rho =0.84 (p<0.01)		
Calhoun et al. 2012 USA Pilot study Shriners Hospitals for Children, Clinical Research Department	N=10 (8M, 2F) Age range: 5-13 years Incomplete: 7 Complete: 3 AIS Score: 3 A, 1 B, 1 C, 5 D	Correlation between WISCI II and SCIM indoor mobility item: r=0.96	Intra-rater reliability: ICC=0.98, CI=0.95-0.99 Inter-rater reliability: ICC=0.97, CI=0.96-0.99	
Burns et al. 2011 USA Test-retest for some participants	N=76 (60M, 16F) Mean age = 43.4±13.8 Mean years from injury = 6.32±5.99 Chronic SCI 45% Paraplegia 55% Tetraplegia	To assess convergent validity for both self-selected and maximum WISCI levels and walking speeds, their relationships with LEMS, UEMS, and MMT were assessed.	ICC for WISCI: SS WISCI – level: 0.994 SS WISCI – speed: 0.930 Max WISCI – level: 0.995	Please see table below.

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
Regional Spinal Cord Injury Center of the Delaware Valley and Magee Rehabilitation Hospital, Philadelphia, PA	<p>Injury etiology: 32% motor vehicle accidents 26% falls 13% sports/diving incidents 11% acts of violence 18% other</p> <p>The distribution of AIS grades was A (3%), B (1%), C (8%), and D (88%), which reflects that participants had to ambulate a minimum of 10 m to be assigned a WISCI level and participate.</p>	<p>For both maximum WISCI and self-selected WISCI, the strongest correlations were with LEMS: $p=0.717$ and $p=0.704$, respectively.</p> <p>There were profound differences when the composite cohort was split into tetraplegic ($n=42$) and paraplegic ($n=34$) cohorts.</p> <p>For tetraplegic participants, there were also significant correlations between WISCI levels and UEMS: $p=0.496$ (self-selected) $p=0.502$ (maximum)</p> <p>Spearman correlations:</p>	Max WISCI – speed: 0.971	

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Btwn Self-selected WISCI level and:</p> <ol style="list-style-type: none"> 1. ASIA UEMS (tetraplegic only, N=41): 0.496 (p<0.0001) 2. ASIA LEMS (N=76): 0.704 (p<0.0001) 3. Manual Muscle Test (Upper & Lower Extremity) (N=75): 0.647 (p<0.0001) <p>Btwn Self-selected WISCI speed and:</p> <ol style="list-style-type: none"> 4. ASIA UEMS (tetraplegic only, N=41): 0.491 (p<0.05) 5. ASIA LEMS (N=76): 0.509 (p<0.05) 6. Manual Muscle Test (Upper & Lower Extremity) (N=75): 0.494 (p<0.0001) 		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Btwn Max WISCI level:</p> <ul style="list-style-type: none"> 7. ASIA UEMS (tetraplegic only, N=41): 0.502 (p<0.0001) 8. ASIA LEMS (N=76): 0.717 (p<0.0001) 9. Manual Muscle Test (Upper & Lower Extremity) (N=75): 0.663 (p<0.0001) <p>Btwn Max WISCI speed:</p> <ul style="list-style-type: none"> 10. ASIA UEMS (tetraplegic only, N=41): 0.469 (p<0.0001) 11. ASIA LEMS (N=76): 0.572 (p<0.0001) 12. Manual Muscle Test (Upper & Lower Extremity) (N=75): 0.539 (p<0.0001) 		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability																		
		<p>More details of paraplegic/tetraplegic values available in article.</p> <p>When the entire cohort was analyzed, walking speed correlated significantly with MMT, LEMS, and WISCI (maximum and self-selected).</p>																				
	<p>SRD for WISCI Level and Walking Speed</p> <table border="1" data-bbox="472 959 1400 1146"> <thead> <tr> <th></th> <th></th> <th>SEM</th> <th>SRD</th> </tr> </thead> <tbody> <tr> <td rowspan="2">SS WISCI</td> <td>Level</td> <td>0.283</td> <td>0.785</td> </tr> <tr> <td>Speed</td> <td>0.091</td> <td>0.254 m/s</td> </tr> <tr> <td rowspan="2">Max WISCI</td> <td>Level</td> <td>0.215</td> <td>0.597</td> </tr> <tr> <td>Speed</td> <td>0.059</td> <td>0.163 m/s</td> </tr> </tbody> </table> <p>WISCI = Walking Index for Spinal Cord Injury</p> <p>SS = Self-Selected</p> <p>Max = Maximum</p>					SEM	SRD	SS WISCI	Level	0.283	0.785	Speed	0.091	0.254 m/s	Max WISCI	Level	0.215	0.597	Speed	0.059	0.163 m/s	
		SEM	SRD																			
SS WISCI	Level	0.283	0.785																			
	Speed	0.091	0.254 m/s																			
Max WISCI	Level	0.215	0.597																			
	Speed	0.059	0.163 m/s																			

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
	SEM = Standard Error of Measurement SRD = Smallest Real Difference			
Burns et al. 2011 USA Validity and reproducibility study Regional Spinal Cord Injury Center of Delaware Valley	N = 76 Mean age: 43.3 ± 13.8 Mean time since injury: 6.32±5.99 years Paraplegic = 34 Tetraplegic = 42 AIS: 3%A, 1%B, 8%C, 88%D	Correlation with ASIA Motor Score: UEMS: =0.496-0.502 (tetraplegic only) LEMS: =0.572-0.717	Test-retest reliability: ICC=0.930-0.995	
Lemay & Nadeau, 2010 Canada Longitudinal An intensive rehabilitation center in	N = 32 SCI (25M, 7F) Mean age: 47.9± 12.8 yrs Neurological level: 15 paraplegic, 17 tetraplegic Level of injury: 17 cervical, 10 thoracic, 5	Spearman's correlations with other walking scales: 1. (all P<0.01) 2. BBS: 0.816 3. SCI-FAI parameter: 0.761 4. SCI-FAI assistive devices: 0.980		Responsiveness: No data available Ceiling effect = 44.8% (44.8% of subjects reached maximal score on the scale)

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>Montreal, Canada (Institut de readaptation Gingras-Lindsay de Montreal)</p>	<p>lumbar Type of injury: 21 traumatic, 11 non- traumatic</p> <p>Inclusion criteria: (1) Adults with SCI AIS D either of traumatic or nontraumatic etiology and (2) the ability to walk 10m independently with or without upper-extremity assistive devices.</p>	<p>5. SCI-FAI mobility: 0.630 6. 2MWT: 0.749 7. 10MWT: 0.795 8. TUG: -0.799</p>		<p>Interpretability: No data available</p>
<p>Marino et al. 2010</p> <p>USA</p> <p>Reliability study</p> <p>Regional Spinal Cord Injury Center of the Delaware Valley</p>	<p>N=26 (9 US, 17 Italy) (16M, 10F)</p> <p>Mean age: 46.4±19.3 years</p> <p>Time post-injury: 8-336 months, mean: 58 months</p> <p>Traumatic SCI = 18</p>		<p>Intra-rater reliability (self-selected (SS), maximum) ICC=1.00</p> <p>Interrater reliability: ICC=1.00 (self selected WISCI) ICC=0.98 (maximum WISCI)</p>	

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
And the the Spinal Unit	Spinal cord lesions = 8 Neurological levels: 7 cervical, 11 thoracic, 8 lumbar AIS: 23D, 2A, 1C			
<p>Marino et al. 2010</p> <p>USA/Italy</p> <p>Reliability study</p> <p>Study subjects were recruited from (1) the Regional Spinal Cord Injury Center of the Delaware Valley, a partnership of Thomas Jefferson University Hospital and</p>	<p>N = 26 SCI (16M, 10F; 9 from USA, 17 from Italy)</p> <p>Mean age: 46.4±19.3 yrs</p> <p>Neurological levels: 7 cervical, 11 thoracic, 8 lumbar</p> <p>AIS A: 2 AIS C: 1 AIS D: 23</p>		<p>Intraclass correlation: coefficients for intrarater reliability were 1.00 for self-selected and maximum WISCI levels for both therapists.</p> <p>Interrater reliability: was 1.00 for self-selected WISCI and 0.98 for maximum WISCI.</p> <p>Bland-Altman plots for differences in time show that the time for the 10-m</p>	

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>Magee Rehabilitation Hospital, Philadelphia, PA, and (2) the Spinal Unit, IRCCS Santa Lucia, Rome, Italy.</p>			<p>walk at SS WISCI varied more from 1 day to the next than between raters on the same day.</p> <p>The difference in time for the two walks on the same day (interrater) was within 25% of the average time in all cases, whereas the difference in time from days 1 to 2 (intrarater) exceeded 25% of average time on several occasions. There was more variability in times for the maximum WISCI than the SS WISCI for both days and raters</p>	
<p>Wirz et al. 2010 Switzerland</p>	<p>N = 42 (33M, 9F) Mean age: 49.3±11.5</p>	<p>WISCI II correlation with: 1. Berg Balance: r=.82 (P<.001)</p>		<p>Responsivness: No data available</p>

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>Prospective study</p> <p>Spinal Cord Injury Center of the Balgrist University Hospital, Zurich, Switzerland</p>	<p>Mean time since injury (SD) = 66.5 months (66.2)</p> <p>AIS A: 2 AIS B: 2 AIS C: 35 AIS D: 3</p> <p>Inclusion criteria: Received either inpatient rehabilitation or out-patient physiotherapy between January 1998 and September 2007. Experienced an SCI at least 1 year prior to enrollment. Able to walk for a minimum distance of 15 m</p>	<p>2. Falls total: $r = -.03$ ($P = .84$)</p> <p>3. SCIM mobility score: $r = .81$ ($P < .001$)</p> <p>4. 10MWT: $r = .81$ ($P < .001$)</p> <p>5. FES-I: $r = -.71$ ($P < .001$)</p> <p>6. Motor scores: $r = .66$ ($P < .001$)</p>		<p>Floor/Ceiling Effect: No data available</p> <p>Interpretability: WISCI mean (SD) score: 16.9 (3.4)</p> <p>Median (range): 18.5 (11-20)</p>
<p>Ditunno et al. 2008</p> <p>Denmark, Germany, Italy, USA</p>	<p>N= 150 (USA = 112; Europe = 38)</p> <p>AIS A: Tetra = 18, Para = 41 AIS B: Tetra = 12, Para = 7</p>	<p>Monotonic Directional Improvement (MDI)</p> <p>77 participants showed improvement, 62/77 participants demonstrated MDI.</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
Prospective cohort	AIS C: Tetra = 22 , Para = 10 AIS D: Tetra = 32, Para = 8	10/15 participants failed to show MDI because a walking device was removed too early. Total Group Spearman correlation w/Lower Extremity Motor Score (LEMS): Initial = 0.47 [P < 0.001] Final = 0.91 [P < 0.001] Improvement = 0.59 [P < 0.0001] Final for those who progressed = 0.71 [P < 0.001] USA Group Spearman correlation w/LEMS: Initial = 0.39 [P < 0.001] Final = 0.91 [P < 0.001] Improvement = 0.54 [P < 0.001]		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Final for those who progressed = 0.79 [P < 0.001]</p> <p>European Group Spearman correlation w/LEMS: Initial = 0.62 [P < 0.001] Final = 0.89 [P < 0.001] Improvement = 0.79 [P < 0.001]</p> <p>Final for those who progressed = 0.42 [P = 0.118]</p> <p>Total Group Spearman correlation w/Locomotor Functional Independence Measure (LFIM): Initial = 0.89 [P < 0.001] Final = 0.76 [P < 0.001] Final for those who progressed = 0.78 [P < 0.001]</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>USA Group Spearman correlation w/LFIM: Initial = 0.89 [P < 0.001] Final = 0.79 [P < 0.001] Final for those who progressed = 0.84 [P < 0.001]</p> <p>European Group Spearman correlation w/LFIM: Final = 0.72 [P < 0.004] Final for those who progressed = 0.72 [P = 0.004]</p>		
<p>Jackson et al. 2008</p> <p>A subcommittee of international experts evaluated locomotion measures</p>	<p>N= 54 expert raters</p>	<p>Content Validity: Expert Evaluations (54 votes): Valid or Useful: 52% Useful but requires validation: 43% Not useful or valid for research: 6%</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>Ditunno et al. 2007</p> <p>USA</p> <p>Single-blinded, paralleled-group, prospective multicenter RCT clinical trial</p> <p>6 regional SCI inpatient rehabilitation centers</p>	<p>N = 146 (114M, 32F)</p> <p>Mean age: 32 (16-69)</p> <p>Level of Injury: 58 cervical, 18 thoracic, 24 lumbar</p> <p>AIS: 36B, 90C, 20D</p> <p>Incomplete spinal cord injury patients who had a Functional Independence Measure locomotor score for walking of < 4 on entry</p>	<p>Correlation with Berg Balance Scale (BBS): r=0.90</p> <p>Correlation with Lower Extremity Motor Score (LEMS): r=0.85</p> <p>Correlation with FIM locomotor score (LFIM): r=0.89</p> <p>Correlation with Functional Independence Measure: r=0.77</p> <p>Correlation with 50-foot walking speed (50FW-S): r=0.85</p> <p>Correlation with 6-minute walking distance (6MW-D): r=0.79</p>		<p>Responsiveness: No data available</p> <p>Interpretability: N=142</p> <p>Mean WISCI (0-20) score: 1.49</p> <p>Floor/ceiling effect At 6 months, the walking speed showed a linear trend to the point of 1 – 1.5 meters/second, and subsequently, a ceiling effect on the WISCI, with walking speed continuing to improve after the WISCI was at or near its maximum value.</p>

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Spearman correlation w/LEMS [all P < 0.001] At 3 months: r = 0.85 At 6 months: r = 0.85 At 12 months: r = 0.88</p> <p>Spearman correlation w/6-Minute Walk Test [all P < 0.001] At 3 months: r = 0.76 At 6 months: r = 0.68 At 12 months: r = 0.69</p> <p>Spearman correlation w/50-foot Walking Speed [all P < 0.001] At 3 months: r = 0.78 At 6 months: r = 0.85 At 12 months: r = 0.77</p> <p>Spearman correlation w/Berg Balance Scale (BBS) [all P < 0.001] At 3 months: r = 0.91</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>At 6 months: $r = 0.89$ At 12 months: $r = 0.92$</p> <p>Spearman correlation w/6-Minute Walk Test [all $P < 0.001$] At 3 months: $r = 0.76$ At 6 months: $r = 0.68$ At 12 months: $r = 0.69$</p> <p>Spearman correlation w/50-foot Walking Speed [all $P < 0.001$] At 3 months: $r = 0.78$ At 6 months: $r = 0.85$ At 12 months: $r = 0.77$</p> <p>Spearman correlation w/Functional Independence Measure (FIM) [all $P < 0.001$] At 3 months: $r = 0.73$ At 6 months: $r = 0.77$ At 12 months: $r = 0.74$</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Spearman correlation w/FIM locomotor score [all P < 0.001] At 3 months: r = 0.92 At 6 months: r = 0.89 At 12 months: r = 0.88</p> <p>Predictors of the WISCI at 12 months (Spearman's rho)</p> <p>Baseline: LEMS = 0.73 BBS = 0.47 FIM Locomotor = 0.30 FIM = 0.12</p> <p>3 Months: LEMS = 0.81 BBS = 0.84 FIM Locomotor = 0.79 FIM = 0.63 Speed = 0.71 Distance = 0.77</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>6 Months: LEMS = 0.86 BBS = 0.89 FIM Locomotor = 0.85 FIM = 0.69 Speed = 0.81 Distance = 0.80</p>		
<p>Kim et al. 2007</p> <p>Prospective cohort</p> <p>Academic medical center.</p>	<p>N = 50 (86%M)</p> <p>Mean age: 47.4 +- 13.2</p> <p>Ambulatory subjects with traumatic incomplete SCI</p>			<p>Reponsiveness: No data available</p> <p>Floor/ceiling effect: Ceiling effect: 48% (24/50) subjects at greater than 1 year post injury has WISCI =20 at entry into the study.</p> <p>Interpretability: No data available</p>
<p>Musselman, 2007</p> <p>Canada</p>	<p>N = 19</p> <p>Incomplete SCI</p> <p>Mean age = 42</p>			<p>1. MCID: 0.06 m/s</p> <p>2. SEM: 0.05 m/s</p> <p>3. Effect Size: 0.46</p>

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>Determining clinical significance via distribution-based and anchor-based approaches</p> <p>Center for Ambulatory Rehabilitation, Research, and Education at the University of Alberta</p>	<p>Time since injury range = 0.6-28.2 years Mean = 6.97 years</p>			
<p>Van Hedel et al. 2006</p> <p>Europe</p> <p>Longitudinal study; analyzed at 1, 3, 6, and 12 months after injury</p>	<p>N = 22 (18M, 4F)</p> <p>Mean age = 45.5±16.7 years (range 17 – 78 years)</p> <p>Incomplete spinal cord injury patients who were able to stand or walk within the first month after SCI.</p>	<p>Spearman correlation w/Lower Extremity Motor Score</p> <p>Within 1 month: r = 0.49 [P=.02]</p> <p>After 3 months: r = 0.50 [P=.02]</p> <p>After 6 months: r = 0.38 [P=.08]</p> <p>After 12 months: r = 0.32 [P=.15]</p>		<p>Responsiveness: 4 time intervals:</p> <p>1) within first month; 2) after 3 months 3) after 6 months; 4) after 12 months:</p> <p>Friedman's test ($\alpha = 0.05$) between 4 intervals: DF = 3, $F_r = 28.7$, $P < 0.001$</p>

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>European Multicenter Study of Human Spinal Cord Injury</p>	<p>Level of Injury: Cervical =13; Thoracic = 1; Lumbar = 7; Sacral = 1</p>	<p>Spearman correlation w/6-Minute Walk Test Within 1 month: r = 0.78 [P<.001] After 3 months: r = 0.28 [P=.20] After 6 months: r = 0.36 [P=.10] After 12 months: r = 0.36 [P=.10]</p> <p>Spearman correlation w/10-Meter Walk Test Within 1 month: r = -0.79 [P<.001] After 3 months: r = -0.21 [P=.35] After 6 months: r = -0.37 [P=.09] After 12 months: r = -0.37 [P=.09]</p>		<p>Pair-wise comparisons via Wilcoxon’s signed rank test:</p> <p>Between intervals 1 and 2: P = 0.005</p> <p>Between intervals 2 and 3: P = 0.18</p> <p>Between intervals 3 and 4: P = 0.31</p> <p>Ceiling effect: All but one of the iSCI subjects qualified up to the max WISCI II score of 20</p> <p>Interpretability: WISCI II mean (SD) score: Within 1st month: 16 (4.6)</p>

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
				After 3 months: 19 (2.4) After 6 months: 20 (0.9) After 12 months: 20 (0.2)
<p>Morganti et al. 2005</p> <p>Italy</p> <p>Retrospective examination</p> <p>Large rehabilitation hospital in center of Italy</p> <p>Rehabilitation hospital in Italy</p>	<p>N=284 (184M, 100F)</p> <p>Mean age: 50.4 ±19.3 (12-86)</p> <p>Mean time post-injury: 56.9±43.9 days</p> <p>Non-traumatic = 177 Traumatic = 107</p> <p>Lesion Level: 81 Cervical, 148 Thoracic, 55 Lumbar-sacral</p> <p>AIS: 84A, 19B, 129C, 52D</p> <p>Concurrent validity sample: N=76</p>	<p>Correlations between:</p> <ol style="list-style-type: none"> 1. WISCI and SCIM: r=0.97 2. WISCI and FIM: r=0.7 3. WISCI and LEMS=0.58 4. WISCI and Barthel Index (BI): r=0.67 5. WISCI and RMI: r=0.67 <p>Groups:</p> <p>Lower Extremity Motor Score (LEM) and WISCI: r=0.58 (p<0.001) (subgroup of 200 patients)</p>	<p>Inter-rater reliability for the WISCI II: r = 1.00 (p<0.001)</p>	

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
	<p>Traumatic or non traumatic SCLs admitted between 1997-2001.</p> <p>Non-traumatic etiology was present in the majority of the patients (177/284): inflammatory (40), vascular (36), neoplastic (39), degenerative (62); traumatic lesions (107/284): car accident (38), motorcycle accident (15), sport accident (7), act of violence (6), suicide attempts (6), and accidental falls (31).</p>	<p>Locomotion outcome at discharge - LEMS and WISCI (eliminating levels 0 and 20): $r=0.57$ ($p<0.001$)</p> <p>Levels at discharge for young patients – LEMS and WISCI: $r=0.50$ ($p<0.01$)</p> <p>Levels at discharge for older patients – LEMS and WISCI: $r=0.64$ ($p<0.01$)</p> <p>Discharge for non-trauma - LEMS and WISCI: $r= 0.58$ ($p<0.01$)</p> <p>Discharge for trauma - LEMS and WISCI: $r= 0.49$ ($p<0.01$)</p> <p>WISCI compared to; Rivermead Mobility Index (RMI): $\rho= 0.67$</p> <p>Barthel Index (BI) $\rho= 0.67$</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Spinal Cord Independence Measure (SCIM): $\rho=0.97$</p> <p>Functional Independence Measure (FIM): $\rho=0.70$</p> <p>RMI and BI: $\rho=0.6$</p> <p>RMI and SCIM: $\rho=0.75$</p> <p>RMI and FIM: $\rho=0.9$</p> <p>BI and SCIM: $\rho=0.7$</p> <p>BI and FIM: $\rho=0.7$</p> <p>SCIM and FIM: $\rho=0.8$</p> <p>All $p < 0.001$</p> <p>WISCI (walking with assistance) levels at discharge and AIS at admission:</p> <p>AIS A vs B: $r=0.573$</p> <p>AIS AB vs C: $r=0.07$</p> <p>AIS AB vs D: $r=0.002$</p> <p>AIS C vs D: $r=0.1$</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>WISCI (independent walking) levels at discharge and ASIA at admission: AIS A vs B: $r=0.02$ AIS AB vs C: $r<0.001$ AIS AB vs D: $r<0.001$ AIS C vs D: $r<0.001$</p> <p>WISCI scale is more sensitive scale for documenting change in levels of walking along a hierarchical order, integrating devices, braces and physical</p>		
<p>Ditunno & Ditunno, 2001</p> <p>USA</p> <p>Retrospective analysis</p> <p>Clinical setting</p>	<p>N=103</p> <p>SCI AIS classification: A=14 B=18 C=52 D=19</p>	<p>Correlation of ASIA grades with WISCI levels were significant: at initial ambulation ($p<0.03$) and at maximum recovery of walking function ($p<0.001$).</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Initial ASIA grades and final WISCI levels correlated at $p < 0.001$.</p> <p>Improvements occurred in one direction in 94% of subjects.</p>		
<p>Ditunno et al. 2000</p> <p>8 SCI centers in Australia, Brazil, Canada, Korea, Italy, the UK, and the USA</p> <p>Methodological study using a modified Delphi technique</p>	<p>N = 24 individuals (8 teams of three composed of health professionals) created this measure.</p>	<p>The WISCI was analyzed to examine whether it appears to measure the construct that it purports to measure. Pilot data at two SCI centers: $W = 0.843$ ($P < .001$)</p> <p>Across all eight SCI centers: International individual data sets: $W = 0.860$ ($P < .001$). Team data sets: $W = 0.872$ ($P < .001$)</p> <p>Sub-group possible pairs of ranking:</p>	<p>100% agreement across all 24 individual international participants and all eight teams.</p>	

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>Clinical physician and Spinal cord injury expert: $\rho=0.968$ ($P<.01$).</p> <p>Physical therapist and Spinal cord injury expert: $\rho=0.944$ ($P<.01$).</p> <p>Physical therapist and Clinical physician: $\rho=0.974$ ($P<.01$)</p> <p>Group Consensus: Using a walker is less impaired than parallel bars.</p> <p>Item 10 was eliminated as there was unacceptable variance.</p> <p>Using a brace, irrespective of one or two canes, reflects a more severely impaired individual than someone without braces.</p>		

Author Year Country Research Design Setting	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		Functional Independence Measure (FIM): $\rho=0.765$ ($P<.001$). 80% of WISCI items fell into two of the FIM categories.		