

Reviewer ID: Emily Procter			
Type of Outcome Measure: Functional Reach Test (FRT) - modified			Total articles: 1
Author ID Year	Study Design	Setting	Population (sample size, age) and Group
Lynch et al. 1998	Test-retest	Magee Rehabilitation Hospital's (Philadel- phia, Pa) SCI follow-up system	N=30 male Mean age 30.8±7.2yrs (range 18-45yrs) AIS A and B (i.e. complete motor) Group 1 – C5-6 (n=10) Group 2 – T1-4 (n=10) Group 3 – T10-12 (n=10)
1. RELIABILITY			
Author ID	Internal Consistency	Test-retest, Inter-rater, Intra-rater	
Lynch et al. 1998	No data available	<i>Mean of three trials taken. This was repeated a second time, 10 minutes following the first test session. There was one rater.</i> Intraclass correlation coefficient. Group 1 – 0.94 Group 2 – 0.85 Group 3 – 0.93	
2. VALIDITY – no data available			
3. RESPONSIVENESS – no data available			
4. FLOOR/CEILING EFFECT – no data available			
5. INTERPRETABILITY			
Author ID	Interpretability		
Lynch et al. 1998	Group 1 (C5-C6) had an SD of 7.6 cm, Group 2 (T1-4) had an SD of 4.3 cm and Group 3 (T10-12) had an SD of 5.6 cm. See top of document for population characteristics. SEM and MDC for mFRT for each group:		
		SEM	MDC
	Group 1 (C5-C6)	1.86	5.16
	Group 2 (T1-T4)	1.67	4.62
	Group 3 (T10-12)	1.48	4.11