

<b>Author Year</b> <b>Country</b> <b>Research Design</b> <b>PEDro Score</b> <b>Total Sample Size</b>	<b>Methods</b>	<b>Outcome</b>
<p><a href="#">Federici et al.</a>, (2019)  Italy  Pre-Post  N=11</p>	<p><b>Population:</b> Mean age: males=50.4±7.3 yr, females=41.5±11.26 yr; Gender: males=5, females=6; Time since injury=30.1±9.4 yr; Level of injury: tetraplegia=3, paraplegia=4, no paraplegia/tetraplegia=4; Severity of injury: complete=5, incomplete=2, none=4.</p> <p><b>Intervention:</b> Participants were 4 couples (one with SCI and one without) and 3 singles with SCI who took part in a sexual health psychoeducational intervention in which 4 couples and 3 singles met every two weeks for 12 meetings of a growth group and reported the results of their love lives and persona lives.</p> <p><b>Outcome Measures:</b> Sexual interest and satisfaction scale (SIS), Beck depression inventory – II (BDI-II) and Beck anxiety inventory (BAI).</p>	<ol style="list-style-type: none"> <li>1. All participants improved significantly on item 5 of the SIS scale “How are your opportunity and your ability to enjoy sexuality yourself?” (p&lt;0.01), SIS scale total score (p&lt;0.05) and BAI scores (p&lt;0.05).</li> <li>2. No difference was found for SIS scale’s general satisfaction after injury, BDI, or BAI (p&gt;0.05).</li> <li>3. Significant effect found on item 5 of the SIS scale “How are your opportunity and your ability to enjoy sexuality yourself?” for both individuals and partners (p&lt;0.05 for both).</li> </ol>