

<b>Author/Year</b>	<b>Treatment (n size)</b>	<b>% of DVTs</b>	<b>Test</b>
Hon et al., (2019)	(n=189)	16.4%	Duplex scan
Morita et al., (2018)	(n=75)	35.7%	Doppler Ultrasonography
Passias et al., (2018)	(n=488,262)	32-36%	
Eichinger et al., (2018)	Prophylactic anticoagulant therapy (n=185)	4%	D-dimer
Clements et al., (2017)	(n=222)	21%	
Marion et al., (2017)	(n=444)	1.6%	
Mackiewicz et al., (2016)	(n=63)	7.9%	D-dimer and venous duplex scans
Piran et al., (2016)	(n=151)	11%	
Dietch et al., (2015)	(n=8,238)	3.5-4.4%	Ultrasonography
Masuda et al., (2015)	(n=268)	10.4%	Ultrasonography
Halim et al., (2014)	LMWH + compression stockings or LMWH only (n=37)	LMWH only=5.4% LMWH + compression stockings=21.6%	Doppler Venous Ultrasonography
Giorgi Pierfranceschi et al., (2013)	LMWH and compression stockings (n=94)	23%	Compression Ultrasound or lower limb colour Doppler ultrasonography, perfusion lung scintigraphy (Q scan) matched with chest X-ray, or computed tomography pulmonary angiography
Germing et al., (2010)	(n=139)	45%	Serial color duplex sonography
Sugimoto et al., (2009)	(n=45)	21%	Doppler Ultrasonography
Colachis & Clinchot (1993)	Prophylaxis Treatment (n=209)	14%	Contrast venography Ultrasound
Gunduz et al., (1993)	Low-dose Heparin (n=31)	53%	Venography
Kulkarni et al., (1992)	Low-dose unfractionated heparin (n=97)	27%	
Merli et al., (1988)	Untreated (n=17)	47%	<sup>125</sup> I fibrinogen scan Impedance Plethysmography Venography
Myllynen et al., (1985)	Anticoagulant therapy (n=37)	100%	<sup>125</sup> I fibrinogen scan Venography
Green et al., (1982)	External pneumatic calf compression (ENCP) or ENCP + aspirin + dipyrid	78% untreated 33% treated	Platelet aggregation studies

	(n=28)		
Rossi et al., (1980)	N/A (n=18)	72%	<sup>125</sup> I fibrinogen scan