

<b>Author Year; Country Score Research Design Total Sample Size</b>	<b>Methods</b>	<b>Outcome</b>
<p><a href="#">Lin et al. 2020</a> China Prospective controlled trial Level 2 N=46</p>	<p><b>Objective:</b> To investigate alterations in fecal microbiome in people with SCI</p> <p><b>Population:</b> SCI group: N=23 Female: 17.3% Age: 32 ± 2.23 years Time since injury: 11 ± 2.68 months Level: 3 cervical 12 thoracic 8 lumbar Severity: 5 complete 18 incomplete Control group: N: 23 Female: 34.8% Age: 28 ± 3.45 years</p> <p><b>Outcome Measures:</b> Microbial communities in the feces of 23 patients and 23 healthy controls were investigated using high-throughput Illumina Miseq sequencing targeting the V3-V4 region of the 16S ribosomal RNA (rRNA) gene. The relative abundances between the fecal microbiota at the genus level in patients with SCI and healthy individuals were determined using cluster analysis.</p>	<ol style="list-style-type: none"> <li>1. People with SCI exhibited microbiome dysbiosis. While there were no significant differences in fecal microbiome alpha diversity (richness and diversity) the structure and quantity significantly differed between SCI and control group (p&lt;0.05)</li> <li>2. 18 operational taxonomic units (OTU) were more abundant in the control group while 27 OTUs were significantly more abundant in the SCI group.</li> </ol>
<p><a href="#">Gungor et al. 2016</a> Prospective</p>	<p><b>Objective:</b> To characterize the gut microbiota in adult</p>	<ol style="list-style-type: none"> <li>1. Results demonstrate that butyrate-producing members</li> </ol>

<p>controlled trial Level 2 N=30</p>	<p>SCI patients with different types of bowel dysfunction. <b>Population:</b> N=40 Level: 15 above T6 in the UMN group 15 with cauda equina syndrome in the LMN group 10 controls Severity: All patients with SCI had ASIA-A injuries UMN group: N=15 Age: 35.0 (9.5) Female: 13.3% Time since injury: 21.0(13.0–105.0) months Level: C4 (n=1, 6.7%) C5 (n=1, 6.7%) C6 (n=1, 6.7%) C7 (n=1, 6.7%) T3 (n=1, 6.7%) T4 (n=4, 26.7%) T5 (n=6, 40.0%) LMN group: N=15 Age: 34.0 (8.9) years Female: 6.7% Time since injury: 18.0(13.0–94.0) months Etiology: 46.7% motor vehicle collisions 26.7% fall 26.7% gunshot wound Level: T12 (n=5, 33.3%) L1 (n=9, 60.0%) L2 (n=1, 6.7%). Control group: Age: 34.4 (8.0) years Female: 0% <b>Outcome Measures:</b> Gut microbial patterns were determined from stool samples</p>	<p>are specifically reduced in SCI patients compared to healthy controls.</p> <ol style="list-style-type: none"> <li>2. Compared to the control group, <i>Pseudobutyrvibrio</i>, <i>Dialister</i> and <i>Megamonas</i> genera were significantly lower in UMN group (<math>p=0.019</math>, <math>p=0.042</math> and <math>p=0.029</math> respectively, Tukey's HSD test) and <i>Roseburia</i>, <i>Pseudobutyrvibrio</i> and <i>Megamonas</i> genera were significantly lower in LMN group (<math>p=0.019</math>, <math>p=0.002</math> and <math>p=0.031</math> respectively, Tukey's HSD test) when compared to healthy controls.</li> <li>3. The <i>Marvinbryantia</i> genus count was significantly lower in UMN bowel dysfunction group (<math>p=0.021</math>, Tukey's HSD test when compared to the LMN group).</li> </ol>
--	--	--

<p><a href="#">Zhang et al. 2018a</a></p> <p>China Prospective Controlled Trial Level 2 N=66</p>	<p><b>Objective:</b> To document neurogenic bowel management of male patients with chronic traumatic complete SCI in the centre, perform comparative analysis of intestinal gut microbiota in male patients with chronic traumatic complete SCI versus males without SCI and explore the association between intestinal microbiota with serum biomarkers and neurogenic bowel symptoms</p> <p><b>Population:</b> 43 SCI 23 non-SCI controls Female: 0% Level: 20 tetraplegia 23 paraplegia Etiology: 37.2% traffic accidents 20.9% bruised by heavy object 20.9% fall from height</p> <p><b>Outcome Measures:</b> microbial diversity by stool sampling; DNA extraction and PCR amplification, Illumina MiSeq sequencing</p>	<ol style="list-style-type: none"> <li>1. Individuals with quadriplegia showed longer time to defecate, higher NBD scores and heavier neurogenic bowel.</li> <li>2. Individuals with quadriplegia showed longer time to defecate, higher NBD scores and heavier neurogenic bowel symptoms than those with paraplegia.</li> <li>3. Gut microbiota diversity in the SCI \group was reduced and had structurally different composition compared to those in the non-SCI adult male group.</li> <li>4. In the SCI group, the abundance of Veillonellaceae and Prevotellaceae increased, while Bacteroidaceae and Bacteroides decreased.</li> <li>5. The abundance of Bacteroidaceae and Bacteroides in the quadriplegia group and Acidaminococcaceae, Blautia, Porphyromonadaceae, and Lachnoclostridium in the paraplegia group were significantly higher than the control group.</li> <li>6. Microbial community structure was significantly associated with serum biomarkers (GLU, HDL, CR, and CRP), NBD defecation time, and COURSE.</li> <li>7. STAMP analysis showed a significant difference (<math>p &lt; 0.05</math>) between the constipation and non-constipation groups (Welch's t-test) in Bifidobacterium on the genus level.</li> <li>8. STAMP analysis showed that Megamonas was significantly higher (<math>p &lt; 0.05</math>) in the bloating group, and Alistipes was significantly higher (<math>p &lt; 0.05</math>) in</li> </ol>
--	--	--

		the without bloating group on the genus level.
<p><a href="#">Yu et al. 2021</a> China Case-control study Level 3 N=69</p>	<p><b>Objective:</b> To explore the hypothesis that 1) the composition and function of gut microbiota are different among patients with complete thoracic SCI, patients with incomplete thoracic SCI and people without SCI and 2) the features of gut microbiota are correlated with the serum biomarkers and implicated in biological functions related to recovery of thoracic SCI</p> <p><b>Population:</b> N=69 Level: 21 Complete thoracic SCI 24 Incomplete thoracic SCI 24 Healthy Etiology: 18 motor vehicle collisions 12 fall from elevated height 7 bruised by heavy object Time since injury: 5.64 ± 2.52 months</p> <p><b>Outcome Measures:</b> NBD score was used to evaluate bowel function for those with SCI. The alpha diversity was determined based on four indices, including observed OTUs (a measure of species richness), Shannon index (a measure of species richness and species evenness), Faith's phylogenetic diversity (a measure of species richness), and Pielou's evenness.</p>	<ol style="list-style-type: none"> <li>1. For the SCI group, there was reduced diversity of the gut microbiota, and alpha diversity had decreased gradually with an increase in the degree of damage.</li> <li>2. Gut microbiota in the SCI group was distinct from non-SCI participants.</li> <li>3. CTSCI group exhibited further deviation in gut microbiota composition than the ITSCI group compared to healthy individuals.</li> <li>4. Four serum biomarkers were found to be correlated with most differential genera.</li> <li>5. SCI accounted for 9.8% (<i>adnois</i> P&lt;0.001) of the gut microbiota variance at the genus level, while the effect size was higher than that observed for other individual. characteristic features, including sex, age, BMI, and clinical serum biomarkers.</li> </ol>
<p><a href="#">Kim et al. 2016a</a> South Korea Pre-post</p>	<p><b>Objective:</b> To investigate the effects and safety of the aqueous extract of the dried,</p>	<ol style="list-style-type: none"> <li>1. Significant decrease in mean (SD) constipation score</li> </ol>

<p>Level 4 N=31</p>	<p>immature fruit of <i>Poncirus trifoliata</i> (L.) Raf., known as <i>Poncirus fructus</i> (PF), in spinal cord injury (SCI) patients with neurogenic bowel.</p> <p><b>Population:</b> N=31 SCI patients with neurogenic bowel (25 were included) Age: 50.9±17.3 years, Range 18-88 years Level: 14 cervical, 11 thoracolumbar, 5 AIS A, 5 AIS B, 4 AIS C, 11 AIS D Etiology: 19 traumatic, 1 transverse myelitis, 5 other Duration of injury: 5.3±6.0 months % Female: 3 females (12%)</p> <p><b>Intervention:</b> <i>Poncirus fructus</i> (PF) administered in dosages of 800 mg each prior to breakfast and lunch for 14 days.</p> <p><b>Outcome Measures:</b> Bowel outcomes before and after administration of PF for 2 weeks. Survey of defecation patterns, plain abdominal radiography, colonic transit times, and side effects</p>	<p>(4.60±3.35 to 3.48±2.42) (p=0.04).</p> <ol style="list-style-type: none"> <li>The Bristol stool scores before and after administration were significantly different (3.52 ± 1.33 to 4.32 ± 1.44 points) (p=0.03).</li> <li>Stool retention score before and after administration of PF was represented with low significance (7.25 +/- 1.60 – 6.46 +/- 1.53 points) in the whole colon (p &lt; 0.05).</li> <li>Colon transit time was significantly in terms of the whole transit time shortened (57.41 ± 20.7 to 41.2 ± 25.5 hours), in right colon (14.4±16.2 to 10.1±12.1h), and in left colon (21.8±12.3 to 14.8±11.8h) (p&lt;0.05).</li> <li>Side effects were observed in 7 people (28.0%) consisting of 2 people with soft stools and 5 people with diarrhea.</li> </ol>
<p><a href="#">Cameron et al. 1996</a> Australia Case Series Level 4 N=11</p>	<p><b>Objective:</b> Assess the nutrient intake of SCI patients, to determine baseline transit time, stool weight and evacuation time and to assess the effect of addition of bran on large bowel function</p> <p><b>Population:</b> Age: range 19-53yrs; Level of injury: C4-T12; 1 participant with incomplete injury and 10 with complete injuries; 7 participants with tetraplegia and 4 with</p>	<ol style="list-style-type: none"> <li>Following the addition of bran, dietary fibre intake significantly increased from 25g/d to 31g/d.</li> <li>Mean colonic transit time significantly lengthened from 28.2 hours to 42.2 hours.</li> </ol>

	<p>paraplegia. All participants were in their first rehabilitation program 1-4 months after injury.</p> <p><b>Treatment:</b> In phase 1 (week 1), participants ate a normal hospital diet and maintained their bowel routine. In phase 2 (week 2-4), fibre intake was increased with the addition of 40g Kellogg's All Bran.</p> <p><b>Outcome Measures:</b> stool weight, total and segmental transit time, bowel evacuation time and dietary intake.</p>	
<b>Author Year; Country Score Research Design Total Sample Size</b>	<b>Methods</b>	<b>Outcome</b>
<p><a href="#">Borsh et al. 2019</a></p> <p>USA Cohort Level 2 N=52</p>	<p><b>Objective:</b> To establish a neurogenic bowel program after SCI in the acute care setting, examine clinician knowledge and ability to deliver the NB program, and evaluate patient knowledge satisfaction and QOL</p> <p><b>Population:</b> N=52 Age at injury: mean 39.3 ± 17.4 y 37% paraplegia 63% tetraplegia Level - 62% cervical 27% thoracic 12% lumbar 40% ASIA A 10% ASIA B 25% ASIA C 13% ASIA D 2% ASIA E 10% unknown</p>	<ol style="list-style-type: none"> <li>1. 77 nurses/patient care technicians and 19 PTs and OTs completed the post-education survey, which reported that knowledge of CPGs improved for all questions after the education in-service.</li> <li>2. Patient knowledge increased significantly from pre-education to post-education including understanding what a SCI is (p=0.02), level of injury (p=0.016), use of suppositories (p=0.008), and digital stimulation (p=0.001).</li> </ol>

	<p><b>Outcome Measures:</b> Pre- and post-education surveys were given to health care providers and patients to measure change in knowledge of NB. Patient survey also included satisfaction and quality of life questions related to health, pain, and self-care taken from the Quality of Life Index-SCI version III. Demographic information, injury characteristics, and bowel medication lists which were extracted from the medical record.</p>	
<p><a href="#">Cabigon et al. 2019</a> USA Cross-sectional Level 5 N=27</p>	<p><b>Objective:</b> The aim of the study was to illustrate how interprofessional collaboration led to utilizing resources of the inpatient rehabilitation facility's peer mentor program and incorporating peer mentors into bowel education for persons with SCI.</p> <p><b>Population:</b> People with SCI who have issues with bowel management N=27 out of 28 responded to the survey</p> <p><b>Outcome Measures:</b> 8-item Likert scale evaluation survey with three open-ended questions to assess the utility of incorporating peer mentors into bowel education</p>	<ol style="list-style-type: none"> <li>1. Results showed that the education program was useful, should be continued, and include the peer mentors.</li> <li>2. Responses from open-ended questions included themes related to knowledge, adherence, and taking charge of one's own care. Interprofessional collaboration and involvement of peer mentors as co-presenters in SCI bowel education were feasible.</li> <li>3. Majority of individuals reported it helped them understand the importance of following a program.</li> </ol>