

Research Summary – Sickness Impact Profile 68 (SIP68) – Quality of Life

Author Year Research Design Setting (country)	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
<p>Post et al. 2001</p> <p>Cross-sectional, using a combination of 3 samples</p> <p>Research lab</p>	<p>N=111 wheelchair- using people (55=SCI and 56=other)</p> <ul style="list-style-type: none"> - Athletes = 49 (SCI = 23, other = 26) - SCI = 32 - Rheumatic disease = 30 <p>Mean age=athletes 36.5, SCI 38.2, Rheumatic disease 69.9</p>	<p>Authors expected moderate to strong correlations between scales of the Nottingham Health Profile (NHP) and the SIP68 with criterion variables, with comparable correlations between scales of each questionnaire that are considered to measure the same aspects of health and the criterion variables.</p> <p>Scales of NHP & SIP68 with Spearman ρ correlations >0.30 (all $P < 0.01$): NHP SA w/ SIP68 Physical Mobility: $\rho = 0.68$</p>	<p>Internal Consistency: $\alpha = 0.88$</p>	

Author Year Research Design Setting (country)	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		NHP ES w/ SIP68 Emotional reactions: $\rho=0.56$ NHP SB w/ SIP68 Emotional reactions: ρ $= 0.41$ NHP PAC w/ SIP68 Emotional reactions: $\rho=0.43$ NHP ES w/ SIP68 Social isolation: $\rho=0.41$ NHP SB w/ SIP68 Social isolation: $\rho=0.35$ NHP PAC w/ SIP68 Social isolation: $\rho=0.43$ NHP ES w/ SIP68 Pain: $\rho=0.36$ NHP SB w/ SIP68 Pain: $\rho=0.54$ NHP MR w/ SIP68 Pain: $\rho=0.38$ NHP PAC w/ SIP68 Pain: $\rho=0.35$ NHP ES w/ SIP68 Energy: $\rho=0.46$		

Author Year Research Design Setting (country)	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>NHP SB w/ SIP68 Energy: $\rho=0.58$ NHP MR w/ SIP68 Energy: $\rho=0.46$ NHP PAC w/ SIP68 Energy: $\rho=0.43$ NHP SB w/ SIP68 Sleep: $\rho=0.37$</p> <p>Abbreviations:</p> <ul style="list-style-type: none">- SA = Somatic Autonomy- MC = Motor Control- ES = Emotional Stability- SB = Social Behaviour- MR = Mobility Range- PAC = Psychologic al Autonomy		

Author Year Research Design Setting (country)	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
		<p>and Communica tion</p> <p>SIP scales & Upper extremity function $\rho=0.13-0.58$ SIP scales & Lower extremity function $\rho=-$ 0.07 to -0.33 SIP scales & Perceived health $\rho=-0.02-0.24$ SIP scales & Social warning $\rho=0.09-0.62$ SIP scales & Well- being $\rho=0.08-0.53$</p>		
<p>Post et al. 1998 (<i>Predictors of Health Status...</i>)</p> <p>Cross-sectional study</p> <p>Community in Netherlands</p>	318 SCI patients; avg. age=39.4; 75% men	<p>Social functioning (path coefficient* = - 0.48), marital status (- 0.38), psychological functioning (-0.19), and age (-0.16) were significant predictors of life satisfaction (total $R^2=0.44$). Authors did not mention if the above</p>		

Author Year Research Design Setting (country)	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability														
		negative correlations were expected. *Path coefficient in multiple regression are comparable to beta values, as calculated by LISREL statistical program																
Post et al. 1996 Cross-sectional, oral interviews Patient's homes	N=315; 75% male, avg. age = 39.4 SCI injury patients between ages 18-65 Complete tetraplegia: 21.7% Complete paraplegia: 29.2% Incomplete tetraplegia: 20.4% Incomplete paraplegia: 28.6% Mean time since injury = 3.6 years, SD 1.9	Spearman correlations: Inter-correlation between $\rho = 0.08-0.67$ on subscales. All correlations are low, except for correlation of "mobility range" with "social behavior" ($\rho = 0.67$). Correlations $\rho>0.20$ (19) have $P<0.001$. Barthel Index and SIP 68, $\rho = -0.74$, $P<.001$ Life Satisfaction Questionnaire (LSQ or LISAT-9) and SIP-68, $\rho = -0.52$, $P<.001$	Internal Consistency: Cronbach's alpha = 0.92 5 out of 6 subscales have good internal consistency 0.72-0.91; The subscale "emotional stability" was the only to lag behind (0.68); but still has 'moderately good' internal consistency.	Interpretability: <table><tr><th>Subscale:</th><th>Mean (SD) score:</th></tr><tr><td>Somatic autonomy</td><td>5.8 (4.8)</td></tr><tr><td>Mobility control</td><td>7.4 (2.4)</td></tr><tr><td>Mobility range</td><td>2.4 (2.4)</td></tr><tr><td>Social behavior</td><td>5.1 (3.1)</td></tr><tr><td>Emotional stability</td><td>1.0 (1.4)</td></tr><tr><td>Psychic autonomy &</td><td>1.1 (1.9)</td></tr></table>	Subscale:	Mean (SD) score:	Somatic autonomy	5.8 (4.8)	Mobility control	7.4 (2.4)	Mobility range	2.4 (2.4)	Social behavior	5.1 (3.1)	Emotional stability	1.0 (1.4)	Psychic autonomy &	1.1 (1.9)
Subscale:	Mean (SD) score:																	
Somatic autonomy	5.8 (4.8)																	
Mobility control	7.4 (2.4)																	
Mobility range	2.4 (2.4)																	
Social behavior	5.1 (3.1)																	
Emotional stability	1.0 (1.4)																	
Psychic autonomy &	1.1 (1.9)																	

Author Year Research Design Setting (country)	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability		
		On Catell's salient similarity index, comparing proposed factor structure to obtained factor structure for all subscales, s values ranged from 0.37-0.95, showing good similarity.		communication		
				Total SIP68	22.8 (11.1)	
Dallmeijer & van der Woude 2001 Cross-sectional study Community in Netherlands	N=37 10 high tetraplegia 9 low tetraplegia 7 motor incomplete tetraplegia 11 paraplegia	Correlation between SIP68 and endurance capacity measures: SIP68 and VO2peak: $r = -0.74$ SIP68 and POmax: $r = -0.68$				
Nanda et al. 2003 Cross-sectional study	N=398, 119 with SCI 49% male Mean age (SD): 53.8 (18.2) years Retest: N=40, all with SCI 100% male	Correlation of SIP68 with: ADL: $r = 0.44$ IADL: $r = 0.57$ SIP: $r = 0.94$	Test-retest, Inter-rater, Intra-rater: Test-retest: ICC = 0.88	Floor/ceiling effect: Ceiling effects are observed for three SIP68 scales: Psychological autonomy and communication: 23.7%		

Reviewer ID: Tyra Chu, Carlos L. Cano-Herrera

Last updated: February 14th, 2024

Author Year Research Design Setting (country)	Demographics and Injury Characteristics of Sample	Validity	Reliability	Responsiveness Interpretability
Multiple communities in the US	Mean age (SD): 51.9 (13.0) years			Emotional stability: 53.6% Mobility range: 23.7%